arXiv:1211.2020v1 [math.CO] 9 Nov 2012

AN UPPER BOUND ON THE COARSENESS OF POINT SETS

J.M. DÍAZ-BÁÑEZ, R. FABILA-MONROY, P. PÉREZ-LANTERO, AND I. VENTURA

Abstract. Let S be a two colored (red and blue) set of n points in the plane. A subset I of S is an island if there is a convex set C such that $I = C \cap S$. The *discrepancy* of an island is the absolute value of the number of red minus the number of blue points it contains. A *convex partition* of S is a partition of S into islands, with disjoint convex hulls. The discrepancy of a convex partition is the discrepancy of its island of minimum discrepancy. The *coarseness* of S is the discrepancy of the convex partition of S with maximum discrepancy. This concept of was recently defined by Bereg et al. [CGTA 2013]. In this paper we prove an almost tight upper bound of $O(n^{1/4}\sqrt{\log n})$ on this parameter. This is obtained by relating the coarseness of a point set to the discrepancy of a certain class of islands.

1. Introduction

Let S be a finite set of n elements, and $\mathcal{Y} \subset 2^S$ a family of subsets of S. The tuple $(S, \mathcal{Y})$ is called a *range space*. If the range space arises from point sets and geometric objects, $(S, \mathcal{Y})$ is called a *geometric range space*. A *coloring* of S is any mapping $\mathcal{X} : S \to \{-1, +1\}$. We think of the elements of S mapped to -1 as being *blue* and the elements of S mapped to $+1$ as being *red*. Let R be the red elements of S and B its blue elements. For $Y \subset S$, let $\mathcal{X}(Y) = \sum_{y \in Y} \mathcal{X}(y)$. The *discrepancy of* $\mathcal{Y}$ is defined as $\operatorname{disc}(\mathcal{Y}) := \min_{\mathcal{X}} \max_{Y \in \mathcal{Y}} |\mathcal{X}(Y)|$.

Geometric discrepancy theory has applications in statistics, clustering, optimization, and computer graphics. See the textbooks [11], [6], [8], [2] and [3] for problems and results in geometric discrepancy.

Assume from now on that S is a set of n points in general position in the plane. A subset I of S is an *island* if there is a convex set C on the plane such that $I = C \cap S$ [4]. A *convex partition* of S is a partition of S into islands, with pairwise disjoint convex hulls. The discrepancy of a convex partition Π of S, $\operatorname{disc}(\Pi)$, is the minimum of $\operatorname{disc}(S_i)$ over $S_i \in \Pi$. The *coarseness* of S is defined as the maximum of $\operatorname{disc}(\Pi)$ over all convex partitions Π of S. This concept was just recently defined in [5]. In this paper we show that the coarseness of S is upper bounded by $O\left(n^{1/4}\sqrt{\log n}\right)$ (Theorem 2.6). We also show that there exists point sets with coarseness at least $\Omega(n^{1/4})$ (Theorem 1.4). We prove the upper bound by showing that the discrepancy of a convex partition is closely related to the discrepancy of a certain class of islands of S, which we call *k-separable* islands.

J.M.Díaz-Bañez and I. Ventura are partially supported by project FEDER MEC MTM2009-08652, and the ESF EUROCORES programme EuroGIGA -ComPoSe IP04-MICINN Project EUI-EURC-2011-4306.

R. Fabila-Monroy is partially supported by CONACyT of Mexico, grant 153984.

P. Pérez-Lantero is partially supported by project FEDER MEC MTM2009-08652 and grant FONDECYT 11110069.

1.1. **Primal and Dual Shatter Functions.**

In this section we recall some definitions and results from discrepancy theory.

The *primal shatter function* $\pi_{\mathcal{Y}}(m)$ of $(S, \mathcal{Y})$ is a function of m. It is defined as the maximum number of subsets into which a subset of S, of at most m elements, can be split (or "shattered") by all the elements of $\mathcal{Y}$. Formally:

$$\pi_{\mathcal{Y}}(m) := \max_{A \subset S, |A| \leq m} |\{Y \cap A : Y \in \mathcal{Y}\}|$$

The *dual shatter function* $\pi^*_{\mathcal{Y}}(m)$ is obtained by exchanging the roles of the points in S with the sets in $\mathcal{Y}$. $\pi^*_{\mathcal{Y}}(m)$ is defined as the maximum number of equivalence classes on S defined by an m-element subfamily $\mathcal{Z} \subset \mathcal{Y}$, where two elements x and y of S are equivalent if they belong to the same sets of $\mathcal{Z}$.

The primal and dual shatter functions have been used to give tight and almost tight upper bounds on the discrepancy of range spaces, via the following theorems (see Chapter 5 of [11]).

Theorem 1.1. **(Primal shatter function bound).** *Let $d > 1$ and C be constants such that $\pi_{\mathcal{Y}}(m) \leq Cm^d$ for all $m \leq n$. Then* $\operatorname{disc}(\mathcal{Y})$ *is upper bounded by* $O\left(n^{1/2-1/2d}\right)$.

Theorem 1.2. **(Dual shatter function bound).** *Let $d > 1$ and C be constants such that $\pi^*_{\mathcal{Y}}(m) \leq Cm^d$ for all $m \leq |\mathcal{Y}|$. Then* $\operatorname{disc}(\mathcal{Y})$ *is upper bounded by* $O\left(n^{1/2-1/2d}\sqrt{\log n}\right)$.

For example, if $\mathcal{H}$ is the family of halfplanes, is easy to see that $\pi_{\mathcal{H}}(m) = O(m^2)$. Thus the discrepancy of halfplanes is $O(n^{1/4})$. It is known that this bound is tight:

Lemma 1.3. ([1, 7]) *For arbitrarily large values of n, there exist sets of n points in general position in the plane such that, given any two coloring of S, a halfplane exists within which one color outnumbers the other by at least $Cn^{1/4}$, for some positive constant C.*

From Lemma 1.3 it is easy to prove the following Theorem.

Theorem 1.4. *For arbitrarily large values of n, there exist sets of n points in general position in the plane with coarseness at least $Cn^{1/4}$ for some positive constant C.*

2. k-SEPARABLE ISLANDS AND CONVEX PARTITIONS

An island I of S is *k-separable* if it can be separated from $S \setminus I$ with at most k halfplanes. That is, if there exists a family $\mathcal{F}$ of at most k halfplanes, such that $I = S \bigcap \mathcal{F}$. We denote the set of k-separable islands with $\mathcal{I}_k$. For constant k, we bound the discrepancy of k-separable islands using its dual shatter function. First we consider points in convex position.

Lemma 2.1. *If k is a positive integer and S a set of n points in convex and general position in the plane then $\pi^*_{\mathcal{I}_k}(m) \leq 4km$.*

Proof. Assume that S is sorted clockwise around its convex hull. Note that any k-separable island must consist of at most k intervals of consecutive points of S in this order. Consider a family of m, k-separable islands. There are at most $2km$ points of S that are the endpoints of any such intervals. There are at most $2km$ regions into which the remaining points (which are not endpoints of any interval) can lie. Thus in total there are at most $4km$ equivalence classes. $\square$

Lemma 2.2. *If k is a positive integer and S a set of n points in general position in the plane then $\pi^*_{\mathcal{I}_k}(m) \leq Cm^2$, for some constant C that depends on k.*

Proof. Let $\mathcal{F}$ be a family of m, k-separable islands on S. We first consider the points lying in the convex hull of some island I of $\mathcal{F}$. Note that the convex hull of I is a set of points in convex position. By Lemma 2.1 these points are in at most $4k(m-1)$ different equivalence classes (when considering the other $m-1$ islands in $\mathcal{F}$). Thus in total there at most $4km^2$ equivalence classes for points in the boundary of some island in $\mathcal{F}$. We now bound the number of equivalence classes for points not lying in the boundary of any island. Each such equivalence class is contained in a cell of the line arrangement defined by the following set of lines $\mathcal{L}$. For each island $I \in \mathcal{F}$, let L_I be the set of at most k lines that separate I from $S \setminus I$. Set $\mathcal{L} := \cup_{I \in \mathcal{F}} L_I$. The line arrangement defined by $\mathcal{L}$ has at most $|\mathcal{L}|^2 = k^2m^2$ cells. By setting $C := k^2 + 4k$ the result follows. $\square$

Using the dual shatter function bound and Lemma 2.2 we obtain:

Theorem 2.3. *Let k be a positive constant and S a set of n points in general position in the plane. The discrepancy of k-separable islands is bounded by $O\left(n^{1/4}\sqrt{\log n}\right)$.*

Note that although k-separable island have small discrepancy this is not the case for islands in general. For example for a two colored set of n points in convex position in the plane there is always an island with discrepancy at least $n/2$. It can be shown that in this case the primal and dual shatter function are equal to 2^m.

We now show that every convex partition must contain a 5-separable island. This follows immediately from:

Lemma 2.4. (Theorem 2 in [9]**)** *A collection of n compact, convex, and pairwise disjoint sets in the plane may be covered with n non-overlapping convex polygons with a total of not more than $6n-9$ sides.*

Theorem 2.5. *Every convex partition Π of S has a 5-separable island.*

We arrive at our main result by combining Theorems 2.5 and 2.3.

Theorem 2.6. *The coarseness of a set of n points in general position in the plane is upper bounded by $O(n^{1/4}\sqrt{\log n})$.*

3. Conclusions

We proved that the discrepancy of k-separable islands is upper bounded by $O(n^{1/4}\sqrt{\log n})$, by showing that its dual shatter function $\pi^*_{\mathcal{I}_k}(m)$ is upper bounded by $O(m^2)$. It is known that the dual shatter function bound can be tight for some range spaces (see [11]). It is not hard to see that the primal shatter function of k-separable islands of point sets in convex position is lower bounded by $\Omega(m^k)$. So the primal shatter function bound can be arbitrarily worse than the dual shatter function bound in this case. It is also interesting to note that the discrepancy of 1-separable islands (or halfspaces) is upper bounded by $O(n^{1/4})$. We leave the exact (asymptotic) computation of the discrepancy of k-separable islands as an open problem.

Using the fact that every convex partition has an island (in this case a 5-separable island) of low discrepancy, we showed that the coarseness of convex partitions is upper bounded by $O(n^{1/4}\sqrt{\log n})$. However Theorem 2.4 provides more information; for any positive constant $c < 1$ there exists a positive integer k_c (depending only on c), so that in every convex partition of S into m islands at least cm of them are c_k-separable (and thus have small discrepancy). We think that computing the exact asymptotic value of the upper bound on the coarseness of point sets is an interesting (and hard) open problem.

4. Acknowledgments

The problems studied here were introduced and partially solved during a visit to University of Valparaiso funded by project Fondecyt 11110069 (Chile).

References

[1] R. Alexander. *Geometric methods in the study of irregularities of distribution.* Combinatorica, 10(2), (1990),pp. 115–136.

[2] J.R. Alexander, J. Beck and W.W.L. Chen. *Geometric discrepancy theory and uniform distribution.* Handbook of Discrete and Computational Geometry (second ed.), CRC Press, (1997), pp. 185–207.

[3] P.K. Agarwal and J. Pach. *Combinatorial Geometry.* Wiley-Interscience Series in Discrete Mathematics and Optimization, 16, (1995), 267–290.

[4] C. Bautista, J.M. Díaz-Báñez, D. Lara, P. Pérez-Lantero, J. Urrutia and I. Ventura. *Computing maximal islands.* Operations Research Letters, 3(39), (2011), pp. 246–251.

[5] S. Bereg, J.M. Díaz-Báñez, D. Lara, P. Pérez-Lantero, C. Seara, J. Urrutia, *On the Coarseness of a Bichromatic Point Set.* Computational Geometry: Theory and Applications, Volume 46, Issue 1, 2013, Pages 65-7, 2012.

[6] B. Chazelle. *The discrepancy method in computational geometry.* Handbook of Discrete and Computational Geometry, CRC Press 44, (2004), pp. 983–996.

[7] B. Chazelle, J. Matoušek and M. Sharir. *An Elementary Approach to Lower Bounds in Geometric Discrepancy.* Discrete Comput Geom 13, (1995), 363–381.

[8] M. Drmota, R.F. Tichy, *Sequences, Discrepancies and Applications*, in: Lecture Notes in Mathematics, vol. 1651, Springer, 1997.

[9] H. Edelsbrunner, A. D. Robison and X. Shen. *Covering convex sets with non-overlapping polygons.* Discrete Mathematics, vol. 81, (1990), pp. 153–164.

[10] J. Matoušek. *Tight upper bounds for the discrepancy of halfspaces.* Discrete & Comput. Geom., 13:593–601, 1995.

[11] J. Matoušek. Geometric Discrepancy: An Illustrated Guide. Springer-Verlag, 1999.

[12] J. Matoušek, E. Welzl, and L. Wernisch.*Discrepancy and e-approximations for bounded VC-dimension.* Combinatorica, 13:455-466, 1993.

Departamento de Matemática Aplicada II, Universidad de Sevilla. Seville, Spain.
E-mail address: `dbanez@us.es`

Departamento de Matemáticas, Cinvestav. Mexico City, Mexico.
E-mail address: `ruyfabila@math.cinvestav.edu.mx`

Escuela de Ingeniería Civil Informática, Universidad de Valparaíso. Valparaiso, Chile.
E-mail address: `pablo.perez@uv.cl`

Departamento de Matemática Aplicada II, Universidad de Sevilla. Seville, Spain.
E-mail address: `iventura@us.es`